

10/564845**IAP20 Rec'd PCT/PTO 13 JAN 2006****Process for Preparing Particles Containing an Antiviral**

5 This invention concerns a process for preparing particles containing antiviral pyrimidines and triazines in the form of a co-precipitate, particles prepared by this process, and pharmaceutical dosage forms comprising these particles.

10 An increasing number of active ingredients having interesting pharmacological activities are hard to formulate into standard oral formulations. This because increased structural complexity or the presence of lipophilic moieties results in a very limited solubility in aqueous media and concomitant reduced or even in some cases absent
15 absorption of the active ingredient in the gastro-intestinal tract. This typically translates into an impaired or a very low bioavailability so that high doses of the active need to be administered to patients in order to achieve effective concentration levels. Therefore a single dose administration may be insufficient for the drug to be effective requiring a multi-dose administration either at once or by a multiple daily dosing, or both.

Since long, formulations have been developed aimed at overcoming these problems. Among these, solid dispersions of active ingredients have been found to be particularly attractive in that they often provide considerable improvement in the solubility and
20 concomitant bioavailability of the active ingredients formulated in this manner. Solid dispersions in general consist of an active ingredient that is dispersed in a hydrophilic medium, usually a polymer.

25 Solid dispersions are usually obtained using two methods. On the one hand they can be obtained via evaporation of a solution consisting of the active ingredient and an inert polymeric material, and on the other via co-fusion of the aforementioned components, with subsequent solidification. In the former instance, the resulting product is referred to as co-precipitate, in the latter as co-melt. Depending on the preparation technique used, co-precipitates can be a solid mass, but they can also be made in particulate form.
30 Solid dispersions in particulate form have become attractive formulation forms for active ingredients with problematic bioavailability.

35 Solid dispersions in particulate form face a number of challenges. First of all they should allow the highest possible level of dissolution and release of the active ingredient in order to ensure sufficient and effective bioavailability thereof.

Secondly, the particles preferably should have a narrow particle size distribution and be of a size that is appropriate for the preparation of dosage forms for oral administration, while containing sufficient high levels of the active ingredient.

Thirdly, the particles should be sufficiently stable over time and able to withstand unfavorable temperature and humidity conditions, such as those commonly used in stability studies, for example storage for extended periods of time at 30°C/60% RH
5 (Relative Humidity), or at 40°C/75% RH.

In certain cases, a solid dispersion in itself is not sufficient to guarantee a sufficient level of bioavailability and of effectiveness of the active ingredient that needs to be administered. To that purpose a number of improvements have been proposed such as
10 the addition of certain ingredients. International patent application WO 97/04749 concerns a preparation process for solid dispersions wherein the active ingredient(s) is/are dissolved in an organic solvent otherwise containing a highly hydrophilic cyclic amide, and advantageously a surface active agent, with the resulting organic solution subsequently evaporated dry, then milled and sieved. The cyclic amide is a
15 polyvinylpyrrolidone with a molecular weight that varies between 10,000 and 50,000.

Certain classes of pyrimidines and triazines have been disclosed in WO-99/50250, WO-99/50256, WO-00/27825 and WO-03/016306 as potent HIV-inhibiting agents. Many of these compounds, in particular those lacking solubilizing groups such as
20 carboxyl or amino groups, have low to very low solubility and need to be formulated in specially adapted formulations in order to provide sufficient bioavailability. Providing formulations for oral dosage forms that deliver an effective amount of these active ingredients poses a particular challenge. This is even more critical given the fact that these active ingredients are HIV-inhibiting agents where it is an absolute requirement
25 that blood plasma levels surpass a certain threshold, which marks the level under which the active no longer is effective amount so that the virus is adequately suppressed thereby avoiding mutations.

WO-01/22938 is concerned with pharmaceutical compositions of pyrimidine and
30 triazine antiviral compounds, in particular the pyrimidines and triazines mentioned above, comprising particles obtainable by melt-extruding a mixture comprising one or more antiviral compounds and one or more appropriate water-soluble polymers and subsequently milling said melt-extruded mixture. A disadvantage associated with melt-extrusion is that the components need to be heated to obtain a homogenous melt so that
35 the active ingredient may be subject to degradation.

The pyrimidine and triazine antivirals mentioned above need to be administered in unit doses containing relatively high amounts of active ingredient, for example amounts which are in the range of 100 mg to 800 mg per unit dose requiring relatively high

amounts of solid dispersion forming polymer. Solid dispersions in particle form containing these active ingredients pose a particular challenge in that a lot of material has to be sprayed on an inert core of restricted size. Indeed, the size of the core is limited because otherwise too much of the core material is present in the end
5 formulation giving rise to unpractically large dosage forms. Moreover particles on which a lot of material has been sprayed on tend to agglomerate and therefore are much more difficult to further process.

The present invention is aimed at overcoming these problems in that it provides
10 processes for preparing formulations comprising particles wherein a co-precipitate is applied as a layer surrounding a neutral hydrophilic core and wherein the co-precipitate comprises, at least one HIV-inhibitory pyrimidine or triazine, one surface-active agent, and one hydrophilic polymer. These particles show improved bioavailability of active ingredients even when almost insoluble in aqueous media, and increase both the
15 dissolution rate and the amount of active ingredient that is released *in vitro* and *in vivo*.

The layer surrounding the neutral hydrophilic carrier is a solid dispersion comprising a co-precipitate, which is soluble in water or in physiological media, such as for example
20 in gastric juice.

The particles prepared according to the process of this invention have the additional advantage of improving the problem of particle agglomeration that is frequently faced when spraying large quantities of coating solutions, by grinding the particles not only after finalization of the spraying step, but also during this step.
25

The size reduction resulting from grinding thus allows to continue spraying on particles of smaller size, that are less likely to agglomerate than particles obtained in a single spraying step with a continuous increase of particle size.

Thus in one aspect, the present invention concerns a process for preparing a particle comprising a co-precipitate surrounding a neutral hydrophilic carrier, said process comprising spraying an organic solution on a neutral hydrophilic carrier, said solution comprising at least one triazine or pyrimidine active ingredient having HIV inhibiting properties, one surface active agent, and one hydrophilic polymer, wherein the spraying
30 of whole of the solution occurs in at least two separate steps, each of these steps followed by a grinding step of the product obtained at the end of the preceding step.

In a particular aspect, the present invention concerns the preparation of particles comprising a co-precipitate surrounding a neutral hydrophilic carrier comprising the following steps:

- 5 a) preparing a solution comprising at least one triazine or pyrimidine active ingredient having HIV inhibiting properties, a hydrophilic polymer, and a surface active agent, in an organic solvent;
- b) spraying part of the solution obtained in step a) on a neutral hydrophilic carrier;
- c) grinding of the particles obtained in step b);
- 10 d) spraying of the remaining quantity of organic solution on the particles obtained in step c) carrier, and
- e) final grinding of the particles obtained in step d).

The spraying/grinding sequence (steps b to d) may be done once, or repeated several
15 times, depending on the volume of the solution intended for spraying and the growth kinetics of the particle during the course of the spraying step.

The carrier used in the process of the invention preferably takes the form of neutral hydrophilic particles, which function as a core on which the solution as specified herein
20 containing the pyrimidine or triazine active ingredient, is sprayed.

All of the organic solution may be prepared in a single step. However, in order to avoid excess evaporation of the organic solvents, it is preferable to prepare each fraction of
25 the solution just prior to the spraying step.

Spraying of the organic solution may be carried out in a conventional fluidized bed drier equipped with a spraying device using the top or bottom spraying process. In a preferred embodiment of the process of the invention, all the spraying steps are carried out in a fluidized bed, equipped with an anti-deflagration device.
30

In certain embodiments, in particular in case of larger batches, the material obtained after grinding (as in step c)), may be partitioned in two or more parts, preferably of equal size. Each part may then be sprayed separately (as in step d)) and ground, whereupon all the resulting material is blended. In alternative embodiments, each of the
35 parts, after spraying (as in step d)) may be ground with a different grinder. The selection of these variants can be done to optimize particle size and particle size distribution.

The fluidized bed is equipped with a spraying nozzle, the position and orientation of which may be selected such that it enables control over the kinetics of the growth of the

particles and avoids problems of sticking, linked to the type of active ingredient, to the qualitative and quantitative composition of the sprayed coating solution, and to various parameters of the procedure (e.g. temperature, air pressure, and the rate of spraying).

- 5 The particles are dried after spraying of the organic solution. Drying may be carried out on trays, or directly inside the equipment used in the spraying step or in a tumble drier. Drying may be carried out either just after spraying of the organic solution, or immediately following the grinding of particles.
- 10 Grinding may be carried out using any type of equipment designed for this particular purpose, and may include use of an oscillating or a blade type grinder. Examples are the FITZMILL rotation type grinder, or the FREWITT oscillating type grinder that are equipped with rotors that force particles through a grid with calibrated openings. The FORPLEX is stator rotor grinder type equipped with pins to which particles are
- 15 projected at a certain feed rate. The type of grinder and the grinding parameters, e.g. speed of grinding, will define the final particle size and particle size distribution.

Different grinders may be used after each spraying step. Moreover, grinding may be done partially on one type of grinder and finished on another type of grinder. All these

20 variants are selected to determine the size and size distribution of the product after grinding.

The process of the invention is particularly useful for active ingredients that are almost insoluble in water, that is for active ingredients where one part is soluble in 1000 parts

25 of water, or more than 1000 parts of water, more in particular for active ingredients that are practically insoluble in water, that is for active ingredients of which one part is soluble in 10,000 parts of water, or more than 10,000 parts of water.

The triazine or pyrimidine active ingredient having HIV inhibiting properties for use in

30 the process of the present invention and in the products obtained therewith preferably is a triazine or pyrimidine derivative that has low solubility in water or aqueous media. Low solubility in this context refers to a compound having a solubility of less than 2 mg/ml or less than 1 mg/ml, or even lower, e.g. less than 0.5 mg/ml or 0.2 mg/ml or even 0.1 mg/ml. Low solubility of active ingredients moreover is meant to include

35 active ingredients that are substantially insoluble, e.g. having a solubility below 0.05 mg/ml or even 0.01 mg/ml. Solubilities are in water or in usual aqueous media that are physiologically relevant (e.g. having a pH between 1 – 8). However, triazines or pyrimidines of higher solubility may be used as well, e.g. having solubilities up to 5 mg/ml or 10 mg/ml or even 20 mg/ml

The triazine or pyrimidine active ingredients for use in the processes and resulting products of this invention in particular are those lacking ionizable groups, e.g. carboxyl or amine groups.


5

Of interest are triazine or pyrimidine active ingredients disclosed in any of WO99/50250, WO99/50256, WO00/27825 or WO03/016306, either specifically or falling under any of the definitions specified in these patent applications, which applications are incorporated herein by reference.

10

The triazine or pyrimidine active ingredient having HIV inhibiting properties for use in the processes and products of this invention in particular is:

(a) an antiviral compound of formula



15 a N-oxide, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof, wherein

Y is CR⁵ or N;

A is CH, CR⁴ or N;

n is 0, 1, 2, 3 or 4;

20 Q is -NR¹R² or when Y is CR⁵ then Q may also be hydrogen;

R¹ and R² are each independently selected from hydrogen, hydroxy, C₁₋₁₂alkyl,

C₁₋₁₂alkyloxy, C₁₋₁₂alkylcarbonyl, C₁₋₁₂alkyloxycarbonyl, aryl, amino, mono- or di(C₁₋₁₂alkyl)amino, mono- or di(C₁₋₁₂alkyl)aminocarbonyl wherein each of the aforementioned C₁₋₁₂alkyl groups may optionally and each individually be

25 substituted with one or two substituents each independently selected from hydroxy, C₁₋₆alkyloxy, hydroxyC₁₋₆alkyloxy, carboxyl, C₁₋₆alkyloxycarbonyl, cyano, amino, imino, aminocarbonyl, aminocarbonylamino, mono- or di(C₁₋₆alkyl)amino, aryl and Het; or

R¹ and R² taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or
30 mono- or di(C₁₋₁₂alkyl)aminoC₁₋₄alkylidene;

R³ is hydrogen, aryl, C₁₋₆alkylcarbonyl, C₁₋₆alkyl, C₁₋₆alkyloxycarbonyl, C₁₋₆alkyl substituted with C₁₋₆alkyloxycarbonyl; and

each R⁴ independently is hydroxy, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, amino carbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, or when Y is CR⁵ then R⁴
35 may also represent C₁₋₆alkyl substituted with cyano or aminocarbonyl;

R⁵ is hydrogen or C₁₋₄alkyl;

L is -X¹-R⁶ or -X²-Alk-R⁷ wherein

R⁶ and R⁷ each independently are phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C₁₋₆alkyl,

- 5 C₁₋₆alkyloxy, C₁₋₆alkylcarbonyl, C₁₋₆alkyloxycarbonyl, formyl, cyano, nitro, amino, and trifluoromethyl; or when Y is CR⁵ then R⁶ and R⁷ may also be selected from phenyl substituted with one, two, three, four or five substituents each independently selected from aminocarbonyl, trihalomethyloxy and trihalomethyl; or when Y is N then R⁶ and R⁷ may also be selected from indanyl or indolyl, each of
10 said indanyl or indolyl may be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylcarbonyl, C₁₋₆alkyloxycarbonyl, formyl, cyano, nitro, amino, and trifluoromethyl; when R⁶ is optionally substituted indanyl or indolyl, it is preferably attached to the remainder of the molecule via the fused phenyl ring. For instance, R⁶
15 is suitably 4-, 5-, 6- or 7-indolyl;

X¹ and X² are each independently -NR³-, -NH-NH-, -N=N-, -O-, -S-, -S(=O)- or -S(=O)₂-;

Alk is C₁₋₄alkanediyl; or

- when Y is CR⁵ then L may also be selected from C₁₋₁₀alkyl, C₃₋₁₀alkenyl, C₃₋₁₀alkynyl,
20 C₃₋₇cycloalkyl, or C₁₋₁₀alkyl substituted with one or two substituents independently selected from C₃₋₇cycloalkyl, indanyl, indolyl and phenyl, wherein said phenyl, indanyl and indolyl may be substituted with one, two, three, four or where possible five substituents each independently selected from halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, C₁₋₆alkyloxycarbonyl, formyl, nitro, amino,
25 trihalomethyl, trihalomethyloxy and C₁₋₆alkylcarbonyl;


aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, nitro and trifluoromethyl;

- Het is an aliphatic or aromatic heterocyclic radical; said aliphatic heterocyclic radical is
30 selected from pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl and tetrahydrothienyl wherein each of said aliphatic heterocyclic radical may optionally be substituted with an oxo group; and said aromatic heterocyclic radical is selected from pyrrolyl, furanyl, thienyl, pyridyl, pyrimidinyl, pyrazinyl and pyridazinyl wherein each of said aromatic heterocyclic radical may
35 optionally be substituted with hydroxy;

or

(b) an antiviral compound of formula

-8-



an *N*-oxide, a pharmaceutically acceptable addition salt, or a stereochemically isomeric forms thereof, wherein

$-b^1=b^2-C(R^{2a})=b^3-b^4=$ represents a bivalent radical of formula

5 $-CH=CH-C(R^{2a})=CH-CH=$ (b-1);

$-N=CH-C(R^{2a})=CH-CH=$ (b-2);

$-CH=N-C(R^{2a})=CH-CH=$ (b-3);

$-N=CH-C(R^{2a})=N-CH=$ (b-4);

$-N=CH-C(R^{2a})=CH-N=$ (b-5);

10 $-CH=N-C(R^{2a})=N-CH=$ (b-6);

$-N=N-C(R^{2a})=CH-CH=$ (b-7);

q is 0, 1, 2; or where possible *q* is 3 or 4;

*R*¹ is hydrogen, aryl, formyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyl, C₁₋₆alkyloxycarbonyl, C₁₋₆alkyl substituted with formyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyloxycarbonyl;

15 *R*^{2a} is cyano, aminocarbonyl, mono- or di(methyl)aminocarbonyl, C₁₋₆alkyl substituted with cyano, aminocarbonyl or mono- or di(methyl)aminocarbonyl, C₂₋₆alkenyl substituted with cyano, or C₂₋₆alkynyl substituted with cyano;

each *R*² independently is hydroxy, halo, C₁₋₆alkyl optionally substituted with cyano or

20 $-C(=O)R^6$, C₃₋₇cycloalkyl, C₂₋₆alkenyl optionally substituted with one or more halogen atoms or cyano, C₂₋₆alkynyl optionally substituted with one or more halogen atoms or cyano, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or di(C₁₋₆alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, $-S(=O)_pR^6$, $-NH-S(=O)_pR^6$, $-C(=O)R^6$, $-NHC(=O)H$, $-C(=O)NHNH_2$, $-NHC(=O)R^6$, $-C(=NH)R^6$ or a radical of formula



25 wherein each A independently is N, CH or CR⁶;

B is NH, O, S or NR⁶;

p is 1 or 2; and

*R*⁶ is methyl, amino, mono- or dimethylamino or polyhalomethyl;


30 L is C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, C₃₋₇cycloalkyl, whereby each of said aliphatic group may be substituted with one or two substituents independently selected from

* C₃₋₇cycloalkyl,

- * indolyl or isoindolyl, each optionally substituted with one, two, three or four substituents each independently selected from halo, C₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, polyhalomethyl, polyhalomethoxy and C₁₋₆alkylcarbonyl,
- 5 * phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl, wherein each of said aromatic rings may optionally be substituted with one, two, three, four or five substituents each independently selected from the substituents defined in R²; or
- L is -X-R³ wherein
- R³ is phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl, wherein each of said
- 10 aromatic rings may optionally be substituted with one, two, three, four or five substituents each independently selected from the substituents defined in R²; and
- X is -NR¹-, -NH-NH-, -N=N-, -O-, -C(=O)-, -CHOH-, -S-, -S(=O)- or -S(=O)₂-;
- Q represents hydrogen, C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl or -NR⁴R⁵; and
- R⁴ and R⁵ are each independently selected from hydrogen, hydroxy, C₁₋₁₂alkyl,
- 15 C₁₋₁₂alkyloxy, C₁₋₁₂alkylcarbonyl, C₁₋₁₂alkyloxycarbonyl, aryl, amino, mono- or di(C₁₋₁₂alkyl)amino, mono- or di(C₁₋₁₂alkyl)aminocarbonyl wherein each of the aforementioned C₁₋₁₂alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy,
- 20 C₁₋₆alkyloxy, hydroxyC₁₋₆alkyloxy, carboxyl, C₁₋₆alkyloxycarbonyl, cyano, amino, imino, mono- or di(C₁₋₆alkyl)amino, polyhalomethyl, polyhalomethoxy, polyhalomethylthio, -S(=O)_pR⁶, -NH-S(=O)_pR⁶, -C(=O)R⁶, -NHC(=O)H, -C(=O)NHNH₂, -NHC(=O)R⁶, -C(=NH)R⁶, aryl and Het; or
- R⁴ and R⁵ taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C₁₋₁₂alkyl)aminoC₁₋₄alkylidene;
- 25 Y represents hydroxy, halo, C₃₋₇cycloalkyl, C₂₋₆alkenyl optionally substituted with one or more halogen atoms, C₂₋₆alkynyl optionally substituted with one or more halogen atoms, C₁₋₆alkyl substituted with cyano or -C(=O)R⁶, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or di(C₁₋₆alkyl)amino, polyhalomethyl, polyhalomethoxy, polyhalomethylthio, -S(=O)_pR⁶, -NH-S(=O)_pR⁶, -C(=O)R⁶, -NHC(=O)H, -C(=O)NHNH₂, -NHC(=O)R⁶, -C(=NH)R⁶ or aryl;
- 30 aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, C₁₋₆alkyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, cyano, nitro, polyhaloC₁₋₆alkyl and polyhaloC₁₋₆alkyloxy;
- Het is an aliphatic or aromatic heterocyclic radical; said aliphatic heterocyclic radical is
- 35 selected from pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl and tetrahydrothienyl wherein each of said aliphatic heterocyclic radical may optionally be substituted with an oxo group; and said aromatic heterocyclic radical is selected from pyrrolyl, furanyl, thienyl, pyridinyl, pyrimidinyl, pyrazinyl and


- pyridazinyl wherein each of said aromatic heterocyclic radical may optionally be substituted with hydroxy; Het is meant to include all the possible isomeric forms of the heterocycles mentioned in the definition of Het, for instance, pyrrolyl also includes 2*H*-pyrrolyl; the Het radical may be attached to the remainder of the molecule of
- 5 formula (I-B) through any ring carbon or heteroatom as appropriate, thus, for example, when the heterocycle is pyridinyl, it may be 2-pyridinyl, 3-pyridinyl or 4-pyridinyl.
- or

(c) a compound of formula



- 10 a *N*-oxide, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof, wherein
- $-a^1=a^2-a^3=a^4-$ represents a bivalent radical of formula
- CH=CH-CH=CH- (a-1);
- N=CH-CH=CH- (a-2);
- 15 $-N=CH-N=CH-$ (a-3);
- N=CH-CH=N- (a-4);
- N=N-CH=CH- (a-5);
- $-b^1=b^2-b^3=b^4-$ represents a bivalent radical of formula
- CH=CH-CH=CH- (b-1);
- 20 $-N=CH-CH=CH-$ (b-2);
- N=CH-N=CH- (b-3);
- N=CH-CH=N- (b-4);
- N=N-CH=CH- (b-5);
- n* is 0, 1, 2, 3 or 4; and in case $-a^1=a^2-a^3=a^4-$ is (a-1), then *n* may also be 5;
- 25 *m* is 1, 2, 3 and in case $-b^1=b^2-b^3=b^4-$ is (b-1), then *m* may also be 4;
- R*¹ is hydrogen; aryl; formyl; C₁-6alkylcarbonyl; C₁-6alkyl; C₁-6alkyloxycarbonyl; C₁-6alkyl substituted with formyl, C₁-6alkylcarbonyl, C₁-6alkyloxycarbonyl, C₁-6alkylcarbonyloxy; C₁-6alkyloxyC₁-6alkylcarbonyl substituted with C₁-6alkyloxycarbonyl;
- 30 each *R*² independently is hydroxy, halo, C₁-6alkyl optionally substituted with cyano or -C(=O)*R*⁶, C₃₋₇cycloalkyl, C₂₋₆alkenyl optionally substituted with one or more halogen atoms or cyano, C₂₋₆alkynyl optionally substituted with one or more halogen atoms or cyano, C₁-6alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or di(C₁₋₆alkyl)amino, polyhalomethyl, polyhalomethylthio, -S(=O)_{*p*}*R*⁶, -NH-S(=O)_{*p*}*R*⁶,

$-C(=O)R^6$, $-NHC(=O)H$, $-C(=O)NHNH_2$, $-NHC(=O)R^6$, $-C(=NH)R^6$ or a radical of formula



wherein each A_1 independently is N, CH or CR^6 ; and

5 A_2 is NH, O, S or NR^6 ;

X_1 is $-NR^5$ -, $-NH-NH$ -, $-N=N$ -, $-O$ -, $-C(=O)$ -, C_{1-4} alkanediyl, $-CHOH$ -, $-S$ -, $-S(=O)_p$ -, $-X_2-C_{1-4}$ alkanediyl- or $-C_{1-4}$ alkanediyl- X_2 -;

X_2 is $-NR^5$ -, $-NH-NH$ -, $-N=N$ -, $-O$ -, $-C(=O)$ -, $-CHOH$ -, $-S$ -, $-S(=O)_p$ -;

10 R^3 is NHR^{13} ; $NR^{13}R^{14}$; $-C(=O)-NHR^{13}$; $-C(=O)-NR^{13}R^{14}$; $-C(=O)-R^{15}$; $-CH=N-NH-C(=O)-R^{16}$; C_{1-6} alkyl substituted with one or more substituents each independently selected from cyano, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ; C_{1-6} alkyl substituted with one or more substituents each independently selected from cyano, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 and wherein 2 hydrogen atoms bound at the same carbon atom are replaced by

15 C_{1-4} alkanediyl; C_{1-6} alkyl substituted with hydroxy and a second substituent selected from cyano, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ; C_{1-6} alkyloxy C_{1-6} alkyl optionally substituted with one or more substituents each independently selected from cyano, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ; C_{2-6} alkenyl substituted with one or more substituents each independently

20 selected from halo, cyano, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ; C_{2-6} alkynyl substituted with one or more substituents each independently selected from halo, cyano, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ; $-C(=N-O-R^8)-C_{1-4}$ alkyl; R^7 or $-X_3-R^7$;

25 X_3 is $-NR^5$ -, $-NH-NH$ -, $-N=N$ -, $-O$ -, $-C(=O)$ -, $-S$ -, $-S(=O)_p$ -, $-X_2-C_{1-4}$ alkanediyl-, $-C_{1-4}$ alkanediyl- X_{2a} -, $-C_{1-4}$ alkanediyl- $X_{2b}-C_{1-4}$ alkanediyl-, $-C(=N-OR^8)-C_{1-4}$ alkanediyl-;

with X_{2a} being $-NH-NH$ -, $-N=N$ -, $-O$ -, $-C(=O)$ -, $-S$ -, $-S(=O)_p$ -; and

with X_{2b} being $-NH-NH$ -, $-N=N$ -, $-C(=O)$ -, $-S$ -, $-S(=O)_p$ -;

30 R^4 is halo, hydroxy, C_{1-6} alkyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, aminocarbonyl, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonyl, formyl, amino, mono- or di(C_{1-4} alkyl)amino or R^7 ;

35 R^5 is hydrogen; aryl; formyl; C_{1-6} alkylcarbonyl; C_{1-6} alkyl; C_{1-6} alkyloxycarbonyl; C_{1-6} alkyl substituted with formyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxycarbonyl or C_{1-6} alkylcarbonyloxy; C_{1-6} alkyloxy C_{1-6} alkylcarbonyl substituted with C_{1-6} alkyloxycarbonyl;

R^6 is C_{1-4} alkyl, amino, mono- or di(C_{1-4} alkyl)amino or polyhalo C_{1-4} alkyl;

- R^7 is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, mono or di(C_{1-6} alkyl)amino C_{1-6} alkyl, formyl, C_{1-6} alkyl-carbonyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, aminocarbonyl, $-CH(=N-O-R^8)$, R^{7a} , $-X_3-R^{7a}$ or $R^{7a}-C_{1-4}$ alkyl;
- R^{7a} is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, mono or di(C_{1-6} alkyl)amino C_{1-6} alkyl, formyl, C_{1-6} alkyl-carbonyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, aminocarbonyl, $-CH(=N-O-R^8)$;
- R^8 is hydrogen, C_{1-4} alkyl, aryl or aryl C_{1-4} alkyl;
- R^9 and R^{10} each independently are hydrogen; hydroxy; C_{1-6} alkyl; C_{1-6} alkyloxy; C_{1-6} alkylcarbonyl; C_{1-6} alkyloxycarbonyl; amino; mono- or di(C_{1-6} alkyl)amino; mono- or di(C_{1-6} alkyl)aminocarbonyl; $-CH(=NR^{11})$ or R^7 , wherein each of the aforementioned C_{1-6} alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C_{1-6} alkyloxy, hydroxy C_{1-6} alkyloxy, carboxyl, C_{1-6} alkyloxycarbonyl, cyano, amino, imino, mono- or di(C_{1-4} alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, $-S(=O)_pR^6$, $-NH-S(=O)_pR^6$, $-C(=O)R^6$, $-NHC(=O)H$, $-C(=O)NHNH_2$, $-NHC(=O)R^6$, $-C(=NH)R^6$, R^7 ; or
- R^9 and R^{10} may be taken together to form a bivalent or trivalent radical of formula
- | | | |
|----|---------------------------------|-------|
| 30 | $-CH_2-CH_2-CH_2-CH_2-$ | (d-1) |
| | $-CH_2-CH_2-CH_2-CH_2-CH_2-$ | (d-2) |
| | $-CH_2-CH_2-O-CH_2-CH_2-$ | (d-3) |
| | $-CH_2-CH_2-S-CH_2-CH_2-$ | (d-4) |
| | $-CH_2-CH_2-NR^{12}-CH_2-CH_2-$ | (d-5) |
| 35 | $-CH_2-CH=CH-CH_2-$ | (d-6) |
| | $=CH-CH=CH-CH=CH-$ | (d-7) |

- R¹¹ is cyano; C₁₋₄alkyl optionally substituted with C₁₋₄alkyloxy, cyano, amino, mono- or di(C₁₋₄alkyl)amino or aminocarbonyl; C₁₋₄alkylcarbonyl; C₁₋₄alkyloxycarbonyl; aminocarbonyl; mono- or di(C₁₋₄alkyl)aminocarbonyl;
- R¹² is hydrogen or C₁₋₄alkyl;
- 5 R¹³ and R¹⁴ each independently are C₁₋₆alkyl optionally substituted with cyano or aminocarbonyl, C₂₋₆alkenyl optionally substituted with cyano or aminocarbonyl, C₂₋₆alkynyl optionally substituted with cyano or aminocarbonyl;
- R¹⁵ is C₁₋₆alkyl substituted with cyano or aminocarbonyl;
- R¹⁶ is C₁₋₆alkyl optionally substituted with cyano or aminocarbonyl, or R⁷;
- 10 p is 1 or 2;
- aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro,
- 15 polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, aminocarbonyl, R⁷ or -X₃-R⁷.

As used in the foregoing definitions and hereinafter halo defines fluoro, chloro, bromo and iodo; polyhalomethyl as a group or part of a group is defined as mono- or polyhalosubstituted methyl, in particular methyl with one or more fluoro atoms, for

20 example, difluoromethyl or trifluoromethyl; polyhaloC₁₋₆alkyl as a group or part of a group is defined as mono- or polyhalosubstituted C₁₋₆alkyl, for example, the groups defined in halomethyl, 1,1-difluoro-ethyl and the like; in case more than one halogen atoms are attached to an alkyl group within the definition of polyhalomethyl or polyhaloC₁₋₆alkyl, they may be the same or different; C₁₋₄alkyl as a group or part of a

25 group encompasses the straight and branched chained saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as, for example, methyl, ethyl, propyl, butyl and the like; C₁₋₆alkyl as a group or part of a group encompasses the straight and branched chained saturated hydrocarbon radicals as defined in C₁₋₄alkyl as well as the higher homologues thereof containing 5 or 6 carbon atoms such as, for example pentyl or

30 hexyl; C₁₋₁₀alkyl as a group or part of a group encompasses the straight and branched chained saturated hydrocarbon radicals as defined in C₁₋₆alkyl as well as the higher homologues thereof containing 7 to 10 carbon atoms such as, for example, heptyl, octyl, nonyl or decyl; C₁₋₁₂alkyl as a group or part of a group encompasses the straight and branched chained saturated hydrocarbon radicals as defined in C₁₋₁₀alkyl

35 as well as the higher homologues thereof containing 11 or 12 carbon atoms such as, for example, undecyl, dodecyl and the like; C₁₋₄alkylidene as a group or part of a group defines bivalent straight and branched chained hydrocarbons having from 1 to 4 carbon atoms such as, for example, methylene, ethylidene, propylidene, butylidene and the

- like; C₁₋₄alkanediyl as a group or part of a group encompasses those radicals defined under C₁₋₄alkylidene as well as other bivalent straight and branched chained hydrocarbons having from 1 to 4 carbon atoms such as, for example, 1,2-ethanediyl, 1,3-propanediyl, 1,4-butanediyl and the like; C₃₋₇cycloalkyl as a group or part of a group is generic to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl; C₃₋₁₀alkenyl as a group or part of a group defines straight and branch chained hydrocarbon radicals containing one double bond and having from 3 to 10 carbon atoms such as, for example, 2-propenyl, 2-butenyl, 2-pentenyl, 3-pentenyl, 3-methyl-2-butenyl, 3-hexenyl, 3-heptenyl, 2-octenyl, 2-nonenyl, 2-decenyl and the like, whereby the carbon atom attached to the pyrimidine ring is preferably an aliphatic carbon atom; C₃₋₁₀alkynyl as a group or part of a group defines straight and branch chained hydrocarbon radicals containing one triple bond and having from 3 to 10 carbon atoms such as, for example, 2-propynyl, 2-butylnyl, 2-pentynyl, 3-pentynyl, 3-methyl-2-butylnyl, 3-hexynyl, 3-heptynyl, 2-octynyl, 2-nonylnyl, 2-decynyl and the like, whereby the carbon atom attached to the pyrimidine ring is preferably an aliphatic carbon atom; C₂₋₆alkenyl defines straight and branched chain hydrocarbon radicals having from 2 to 6 carbon atoms containing a double bond such as ethenyl, propenyl, butenyl, pentenyl, hexenyl and the like; C₂₋₁₀alkenyl defines straight and branched chain hydrocarbon radicals having from 2 to 10 carbon atoms containing a double bond such as the groups defined for C₂₋₆alkenyl and heptenyl, octenyl, nonenyl, decenyl and the like; C₂₋₆alkynyl defines straight and branched chain hydrocarbon radicals having from 2 to 6 carbon atoms containing a triple bond such as ethynyl, propynyl, butynyl, pentynyl, hexynyl and the like; C₂₋₁₀alkynyl defines straight and branched chain hydrocarbon radicals having from 2 to 10 carbon atoms containing a triple bond such as the groups defined for C₂₋₆alkynyl and heptynyl, octynyl, nonynyl, decynyl and the like; C₁₋₃alkyl as a group or part of a group encompasses the straight and branched chain saturated hydrocarbon radicals having from 1 to 3 carbon atoms such as, methyl, ethyl and propyl; C₄₋₁₀alkyl encompasses the straight and branched chain saturated hydrocarbon radicals as defined above, having from 4 to 10 carbon atoms. The term C₁₋₆alkyloxy defines straight or branched chain saturated hydrocarbon radicals such as methoxy, ethoxy, propyloxy, butyloxy, pentyloxy, hexyloxy, 1-methylethyloxy, 2-methylpropyloxy, 2-methylbutyloxy and the like; C₃₋₆cycloalkyloxy is generic to cyclopropyloxy, cyclobutyloxy, cyclopentyloxy and cyclohexyloxy.
- As used herein before, the term (=O) forms a carbonyl moiety when attached to a carbon atom, a sulfoxide group when attached once to a sulfur atom, and a sulfonyl group when attached twice to a sulfur atom.

When any variable (e.g. aryl, R^3 , R^4 in formula (I-A) etc.) occurs more than one time in any constituent, each definition is independent.

5 Lines drawn into ring systems from substituents indicate that the bond may be attached to any of the suitable ring atoms. For instance for compounds of formula (I-A), R^4 can be attached to any available carbon atom of the phenyl or pyridyl ring.

The addition salts as mentioned herein are meant to comprise the therapeutically active addition salt forms which the compounds of formula (I-A), (I-B) or (I-C) are able to
10 form with appropriate acids, such as, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid; sulfuric; nitric; phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, methane-sulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic,
15 *p*-aminosalicylic, pamoic and the like acids.

The pharmaceutically acceptable addition salts as mentioned hereinabove are also meant to comprise the therapeutically active non-toxic base, in particular, a metal or amine addition salt forms which the compounds of the present invention are able to
20 form. Said salts can conveniently be obtained by treating the compounds of the present invention containing acidic hydrogen atoms with appropriate organic and inorganic bases such as, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, *N*-methyl-D-glucamine, hydrabamine salts, and
25 salts with amino acids such as, for example, arginine, lysine and the like. Conversely said salt forms can be converted by treatment with an appropriate base or acid into the free acid or base form.

The term addition salts also comprises the hydrates and the solvent addition forms
30 which the compounds of formula (I-A), (I-B) or (I-C) are able to form. Examples of such forms are e.g. hydrates, alcoholates and the like.

Interesting compounds of formula (I-A) are those wherein Y is CR^5 or N; A is CH , CR^4 or N; n is 0, 1, 2, 3 or 4; Q is $-NR^1R^2$; R^1 and R^2 are each independently selected from
35 hydrogen, hydroxy, C_{1-12} alkyl, C_{1-12} alkyloxy, C_{1-12} alkylcarbonyl, C_{1-12} alkyloxy-carbonyl, aryl, amino, mono- or di(C_{1-12} alkyl)amino, mono- or di(C_{1-12} alkyl)amino-carbonyl wherein each of the aforementioned C_{1-12} alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C_{1-6} alkyloxy, hydroxy C_{1-6} alkyloxy, carboxyl, C_{1-6} alkyloxy-

- carbonyl, cyano, amino, imino, aminocarbonyl, aminocarbonylamino, mono- or di(C₁₋₆alkyl)amino, aryl and Het; or R¹ and R² taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C₁₋₁₂alkyl)aminoC₁₋₄alkylidene; R³ is hydrogen, aryl, C₁₋₆alkylcarbonyl, C₁₋₆alkyl, C₁₋₆alkyloxycarbonyl, C₁₋₆alkyl
- 5 substituted with C₁₋₆alkyloxycarbonyl; each R⁴ independently is hydroxy, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethoxy; R⁵ is hydrogen or C₁₋₄alkyl; L is -X¹-R⁶ or -X²-Alk-R⁷ wherein R⁶ and R⁷ each independently are phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy,
- 10 C₁₋₆alkylcarbonyl, C₁₋₆alkyloxycarbonyl, formyl, cyano, nitro, amino, and trifluoromethyl, X¹ and X² are each independently -NR³-, -NH-NH-, -N=N-, -O-, -S-, -S(=O)- or -S(=O)₂-, and Alk is C₁₋₄alkanediyl; aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, nitro and trifluoromethyl; Het is an aliphatic or
- 15 aromatic heterocyclic radical; said aliphatic heterocyclic radical is selected from pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl and tetrahydrothienyl wherein each of said aliphatic heterocyclic radical may optionally be substituted with an oxo group; and said aromatic heterocyclic radical is selected from pyrrolyl, furanyl, thienyl, pyridyl, pyrimidinyl, pyrazinyl and pyridazinyl wherein
- 20 each of said aromatic heterocyclic radical may optionally be substituted with hydroxy.

Preferred compounds of formula (I-A) are


- 4-[[4-amino-6-[(2,6-dichlorophenyl)methyl]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 25 4-[[4-[(2,6-dimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-(2,4,6-trimethylphenoxy)-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[(2,6-dichlorophenyl)thio]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[(2,4-dichloro-6-methylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[2-[(cyanophenyl)amino]-4-pyrimidinyl]amino]-3,5-dimethylbenzonitrile;
- 30 4-[[4-[(2,4-dibromo-6-fluorophenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[(2,4,6-trichlorophenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[2-amino-6-[(2,4,6-trimethylphenyl)amino]-4-pyrimidinyl]amino]benzonitrile;
- 4-[[2-[(4-cyanophenyl)amino]-5-methyl-4-pyrimidinyl]amino]-3,5-dimethylbenzonitrile;
- 35 [[2-[(4-cyanophenyl)amino]-5-methyl-4-pyrimidinyl]amino]-3,5-dimethylbenzonitrile;
- 4-[[4-amino-6-[(2,4,6-trimethylphenyl)amino]-1,3,5-triazin-2-yl]amino]benzonitrile;
- 4-[[4-(hydroxyamino)-6-[(2,4,6-trimethylphenyl)amino]-1,3,5-triazin-2-yl]amino]-benzonitrile;

-17-

- 4-[[4-(hydroxyamino)-6-[(2,4,6-trichlorophenyl)amino]-1,3,5-triazin-2-yl]amino]-benzonitrile;
 4-[[4-(4-acetyl-2,6-dimethylphenoxy)-6-amino-1,3,5-triazin-2-yl]amino]benzonitrile;
 [[4-[(5-acetyl-2,3-dihydro-7-methyl-1H-inden-4-yl)oxy]-6-amino-1,3,5-triazin-2-yl]-
 5 amino]benzonitrile;
 4-[[4-amino-6-[(2-chloro-4,6-dimethylphenyl)amino]-1,3,5-triazin-2-yl]amino]-benzonitrile;
 4-[[4-amino-6-[(2,6-dibromo-4-methylphenyl)amino]-1,3,5-triazin-2-yl]amino]-benzonitrile; the *N*-oxides, the pharmaceutically acceptable addition salts and the
 10 stereochemically isomeric forms thereof.

Most preferred is

- 4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile, or a pharmaceutically acceptable salt thereof, also referred to as TMC120, which can be
 15 represented by the following chemical structure:



- Interesting compounds of formula (I-B) are those wherein one or more of the following
 20 restrictions apply :
- i) $-b^1=b^2-C(R^{2a})=b^3-b^4=$ is a radical of formula (b-1);
 - ii) q is 0;
 - iii) R^{2a} is cyano or $-C(=O)NH_2$, preferably R^{2a} is cyano;
 - iv) Y is cyano, $-C(=O)NH_2$ or a halogen, preferably a halogen;
 - 25 v) Q is hydrogen or $-NR^4R^5$ wherein R^4 and R^5 are preferably hydrogen;
 - vi) L is $-X-R^3$ wherein X is preferably NR^1 , O or S, most preferably X is NH, and R^3 is substituted phenyl with C_{1-6} alkyl, halogen and cyano as preferred substituents.

- Another interesting group of compounds of formula (I-B) are those compounds of
 30 formula (I-B) wherein L is $-X-R^3$ wherein R^3 is 2,4,6-trisubstituted phenyl, each substituent independently selected from chloro, bromo, fluoro, cyano or C_{1-4} alkyl.

Also interesting are those compounds of formula (I-B) wherein Y is chloro or bromo and Q is hydrogen or amino.


Particular compounds of formula (I-B) are those compounds of formula (I-B) wherein
5 the moiety in the 2 position of the pyrimidine ring is a 4-cyano-anilino group.

Further particular compounds of formula (I-B) are those compounds of formula (I-B)
wherein the moiety in the 2 position of the pyrimidine ring is a 4-cyano-anilino group,
L is -X-R³ wherein R³ is a 2,4,6-trisubstituted phenyl, Y is a halogen and Q is hydrogen
10 or NH₂.


Most preferred compounds of formula (I-B) are :

4-[[4-amino-5-chloro-6-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]-
benzonitrile;
15 4-[[5-chloro-4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
4-[[5-bromo-4-(4-cyano-2,6-dimethylphenoxy)-2-pyrimidinyl]amino]benzonitrile;
4-[[4-amino-5-chloro-6-[(4-cyano-2,6-dimethylphenyl)amino]-2-pyrimidinyl]amino]-
benzonitrile;
4-[[5-bromo-6-[(4-cyano-2,6-dimethylphenyl)amino]-2-pyrimidinyl]amino]-
20 benzonitrile;
4-[[4-amino-5-chloro-6-(4-cyano-2,6-dimethylphenoxy)-2-pyrimidinyl]amino]-
benzonitrile; and
4-[[4-amino-5-bromo-6-(4-cyano-2,6-dimethylphenoxy)-2-pyrimidinyl]amino]-
benzonitrile; the *N*-oxides, the pharmaceutically acceptable addition salts, quaternary
25 amines and the stereochemically isomeric forms thereof.

The most preferred compound of formula (I-B) is
4-[[4-amino-5-bromo-6-(4-cyano-2,6-dimethylphenoxy)-2-pyrimidinyl]amino]-
benzonitrile; or a pharmaceutically acceptable addition salt thereof,
30 which usually is designated as TMC125 and which can be represented by formula:



Particular compounds of formula (I-C) are those having the formula



5 wherein

R^1 , R^2 , R^3 , R^4 and X_1 are as defined above;

n' is 0, 1, 2 or 3 ;

$R^{2'}$ is halo, C_{1-6} alkyl, trihalomethyl, trihalomethyloxy, cyano, aminocarbonyl, C_{1-6} alkyl substituted with cyano or aminocarbonyl;

10 and $R^{2'}$ is positioned in the para position in respect of the NR^1 moiety.

Further particular compounds of formula (I-C) are those having the formula (I-C-1) wherein $R^{2'}$ is cyano, aminocarbonyl or C_{1-6} alkyl substituted with cyano or aminocarbonyl.

15

Further particular compounds are those compounds of formula (I-C) or (I-C-1) wherein one or more, preferably all of the following restrictions apply :

a) n is at least 1, in particular 1; or n' is 0;

b) R^2 or $R^{2'}$ is cyano;

20 c) m is 1, 2 or 3;

d) R^4 is C_{1-6} alkyl, especially methyl; nitro; amino; halo; C_{1-6} alkyloxy or R^7 ;

e) R^3 is R^7 , $NR^{13}R^{14}$, $-C(=O)R^{15}$, $-CH=N-NH-C(=O)R^{16}$, $-C(=O)NHR^{13}$,

$-C(=O)NR^{13}R^{14}$, $-C(=N-OR^8)-C_{1-4}$ alkyl, C_{1-6} alkyl substituted with cyano, C_{1-6} alkyl substituted twice with cyano, C_{1-6} alkyl substituted with NR^9R^{10} , C_{1-6} alkyl substituted with hydroxy and cyano, C_{1-6} alkyl substituted with hydroxy and R^7 , C_{1-6} alkyloxy C_{1-6} alkyl, C_{1-6} alkyloxy C_{1-6} alkyl substituted with cyano, C_{2-6} alkenyl substituted with R^7 , C_{2-6} alkenyl substituted with cyano, C_{2-6} alkenyl substituted twice with cyano, C_{2-6} alkenyl substituted with cyano and R^7 , C_{2-6} alkenyl substituted with cyano and

25

-20-


-C(=O)-C₁₋₆alkyl, C₂₋₆alkenyl substituted with cyano and halo, C₂₋₆alkenyl substituted with -C(=O)-NR⁹R¹⁰, C₂₋₆alkenyl substituted with halo, C₂₋₆alkenyl substituted twice with halo or C₂₋₆alkenyl substituted with NR⁹R¹⁰;

f) X₃ is -C(=O)-, -CH₂-C(=O)-, or -C(=N-OR⁸)-C₁₋₄alkanediy-;

5 g) X₁ is NH or O;


h) R¹ is hydrogen or C₁₋₄alkyl.

Preferred compounds of formula (I-C) are:




10

Comp No.	R ³	R ⁴
1	-CH=CH-CN	H
2	-C(CH ₃)=CH-CN	H
3	-CH=C(CH ₃)-CN	H



Comp. No.	R ³	R ¹
4	-CH=CH-CN	H
5	-CH=C(CH ₃)-CN	H



15

Comp. No.	R ³	R ^{4a}	R ^{4b}	X ¹
6	-CH=C(CH ₃)-CN	CH ₃	Cl	-NH
7	-CH=CH-CN	CH ₃	2-furanyl	-NH
8	-CH=C(CH ₃)-CN	CH ₃	Br	-NH
9	-CH=CH-CN	CH ₃	Br	-NH
10	-CH=CH-CN	CH ₃	Cl	-NH

Of particular interest is compound 1 listed above, or a pharmaceutically acceptable salt thereof, which compound is also referred to as TMC278.

5 The active ingredient may be present in the particle in an amount that may vary from 1 to 60% by weight, in particular from 5 to 50% by weight, more in particular from 10 to 40% or from 10 to 30% by weight.

10 The inert hydrophilic carrier may consist of any chemically and pharmaceutically inert excipient, existing in particle form, crystalline or amorphous, for example sugars or sugar derivatives such as lactose, preferably Extra Fine Lactose (EFK), sucrose, hydrolyzed starch (maltodextrins), celluloses, or mixtures thereof such as sucrose and starch; or mixtures with a cellulose base may also be used for the preparation of the inert carrier.

15 The inert hydrophilic carrier is present in an amount that may be as high as 95 % by weight, in particular from 1-95% by weight, more in particular from 5-50% by weight, further in particular from 10 to 40% or from 10-30% by weight.

20 The unit particle size of the inert hydrophilic carrier may range from 50 μm to 500 μm , preferably from 90 μm to 200 μm . Preferred is a carrier that is spherically shaped.

25 Suitable hydrophilic polymers are polyvinylpyrrolidones, cellulose derivatives, acrylic polymers or polyethylene glycols. Of particular interest is polyvinylpyrrolidone, in particular the latter polymer having a molecular weight ranging from 10,000 to 50,000.

30 Appropriate cellulose derivatives comprise hydroxyalkylated celluloses, for example, hydroxypropylmethylcellulose, hydroxypropylcellulose, hydroxymethylcellulose, hydroxypropylmethylcellulose phthalate and hydroxypropylmethylcellulose aceto-succinate.

35 Preferred is hydroxypropylmethylcellulose (HPMC), more in particular the latter polymer having an apparent viscosity ranging from 2.4 to 18 cP, and even more preferably ranging from 2.4 to 5 cP.

40 The acrylic polymer may be selected among ammonium methacrylate copolymer, polyacrylate, polymethacrylate, and methacrylic acid copolymer.

The polyethylene glycol may be chosen among polymers with a molecular weight ranging from 1,000 to 20,000.

The weight ratio of the hydrophilic polymer to the active ingredient ranges preferably from 10:1 to 1:2.

5 The surface-active agent may be selected among cationic, anionic, non-ionic or amphoteric surfactants, separately, or as a mixture.

The surface-active agent may be selected among compounds such as sodium laurylsulfate; the monooleate, monolaurate, monopalmitate, monostearate, trioleate, tristearate, or any other polyoxyethylenated sorbitan fatty ester, preferably Tween® 20,
10 40, 60 or 80; ethoxylated fatty acid glycerides, said fatty acids being saturated or not, and containing at least 8 carbon atoms; poloxamers such as poloxamer 188; bloc copolymers of ethylene oxide/propylene oxide such as Pluronic® F68 or F87; stearyl alcohol, cetostearyl alcohol; lecithin; cholesterol; polyethoxylated castor oil; polyethoxylated fatty alcohol ethers, such as Brij®; and polyethoxylated stearates.

15 The surface-active agent is advantageously present in amounts that may vary between 0.1 and 20% by weight, in particular between 1-10% by weight, more in particular between 1-5% by weight, compared to the total resulting mass.

20 The organic solvent may be selected among an alcohol such as ethanol, isopropanol; an ether such as tetrahydrofuran, isopropyl ether; a ketone such as acetone, methylethylketone; methylene chloride; or any mixture thereof.

25 The amount of solvent used should be selected such that it takes into account the solubility of the various components of the organic solution.

In a further aspect the present invention also concerns a particle comprising, or in an alternative aspect, consisting of, a co-precipitate applied in a layer surrounding a neutral hydrophilic carrier and comprising at least one antiviral pyrimidine or triazine,
30 at least one surface-active agent, and at least one hydrophilic polymer. The neutral hydrophilic carrier preferably is present as a core.

In still a further aspect the present invention also concerns a particle containing or in an alternative aspect, consisting of a co-precipitate applied in a layer surrounding a neutral
35 hydrophilic carrier and comprising at least one antiviral pyrimidine or triazine, at least one surface-active agent, and at least one hydrophilic polymer, obtainable or obtained by the process described herein.

The size of particles usually ranges from 50 μm to 1000 μm , preferably from 100 μm to 800 μm , more preferably from 150 μm to 300 μm , e.g. about 250 μm , and is determined using conventional methods, for example with a set of sieves containing calibrated openings, or using laser diffraction.

5

In a preferred embodiment of the invention, the particles of the invention contain:

- from 15 to 40 % in weight of an inert hydrophilic carrier, preferably lactose, more preferably lactose EFK;
- 10 - from 15 to 30% in weight of an active ingredient as defined herein, which active ingredient preferably is TMC120, TMC125, or TMC278.
- from 30 to 60% in weight of a hydrophilic polymer, preferably HPMC;
- from 1 to 10% in weight of a surface-active agent, preferably a non-ionic surfactant such as polysorbate, in particular selected from the group comprising polysorbates 20 to 80.

15

The particles according to the invention may be used directly, or in a mixture, with excipients typically used in the pharmaceutical art for the preparation of a pharmaceutical form intended for oral administration, such as for example, a capsule, or a tablet, chemically compatible with the active ingredient(s).

20

Thus in still a further aspect, this invention provides a pharmaceutical composition comprising particles as specified herein and one or more further excipients. The invention also provides an oral dosage form comprising particles as specified herein, or comprising a pharmaceutical composition comprising particles as specified herein.

25

Examples of excipients comprise diluents, disintegrating agents, bulking agents, lubricants, antistatic agents, binders or any other adjuvants used in preparing pharmaceutical formulations.

30

The diluent may be selected among sugars such as sucrose, lactose, fructose, dextrose, or polyols with less than 13 atoms of the carbon, such as mannitol, xylitol, sorbitol, maltitol, lactitol, erythritol, dicalcium phosphate, tricalcium phosphate, or microcrystalline cellulose, separately or in combination.

35

The diluent is used in an amount ranging from 20 to 90% in weight, preferably from 30 to 60% in weight, calculated based on the weight of the tablet.

The diluent is preferably used in a directly compressible form, with an average particle diameter ranging from 100 μm to 500 μm , or in the form of a powder with an average particle size that is less than 100 μm , said powder being used separately, or in a mixture with the directly compressible product.

5

The disintegrating agent is selected from the group comprising in particular cross-linked sodium carboxymethylcellulose designated by the term croscarmellose, cross-linked polyvinyl pyrrolidones designated by the term crospovidone, and mixtures thereof.

10

The disintegrating agent is used in an amount ranging from 1 to 20% in weight, preferably from 5 to 15% in weight, calculated based on the weight of the tablet.

The bulking agent is selected from the group comprising microcrystalline cellulose, starches, modified starches, such as carboxymethyl starch or sodium glycolated starch, alginic acid, or sodium alginate, and mixtures thereof.

15

The bulking agent is used in an amount ranging from 1 to 15 % in weight, calculated on the basis of the tablet weight.

20

The lubricant is selected from the group comprising magnesium stearate, stearic acid, sodium stearyl fumarate, polyoxyethylene glycols, sodium benzoate, a pharmaceutically acceptable oil, preferably dimethicone, paraffin oil, or cotton seed oils, and mixtures thereof.

25

The lubricant is used in an amount of up to 2%, preferably from 0.02 to 2% in weight, more preferably from 0.5 to 1% in weight, calculated based on the weight of the tablet.

According to a first variant of preparing the dosage forms of this invention, the total amount of lubricant is incorporated in the mixture of excipients to be compressed; in a second variant, at least a fraction of the lubricant is sprayed on the walls of the matrix and the punches at the moment of compression; the lubricant is then in the form of a powder, for example magnesium stearate or a liquid, for example liquid paraffin oil.

30

The quantities of lubricant used during the internal and/or external steps are carefully adjusted so as to avoid excess that would interfere with cohesion of the powder bed during compression.

35

-25-

The antistatic agent may be selected from the group comprising micronized or non-micronized talc, colloidal silica (e.g. Aerosil®200), processed silica (e.g. Aerosil®R972), or precipitated silica (e.g. Syloid®FP244), and mixtures thereof.

- 5 The antistatic agent is used in an amount of up to 5% in weight, calculated based on the weight of the tablet.

The binder is used in dry form, and may consist of starch, sugar, polyvinylpyrrolidone, or carboxymethylcellulose, separately or as a mixture.

10

The binder is used in an amount of up to 15% in weight, preferably less than 10% in weight, calculated based on the weight of the tablet.

- 15 Adjuvants may also be added to the mixture formulated for capsules, or for compression, and may be selected from pH adjusting agents, systems allowing to produce effervescence, in particular carbon dioxide generators of the type used as pH adjusters, or surface active agents.

- 20 The invention also concerns pharmaceutical dosage forms comprising the particles as described herein.

Preparation of pharmaceutical dosage forms comprising the invention particles may consist of the following steps:

- 25 - dry mixing of the particles obtained according to the procedures previously described, with the excipients;
- compression of the mixture or transfer of the mixture to capsules to obtain a unit form.

- 30 In order to avoid sticking, the punches and/or the internal walls of the compression matrix (the dies) may be treated with a suitable lubricant such as any of the lubricants mentioned above.

- 35 According to one embodiment, the preparation of the dosage forms comprises first mixing the active ingredient and all of the other excipients, including the lubricant whereupon the mixture is compressed. In another embodiment, only part of the lubricant is added, the remainder being sprayed on the punches and/or dies. In still another embodiment, the preparation of the dosage forms comprises two steps, first mixing the active ingredient with all of the compression excipients except for the
40 internal lubricant, and then in a second step, the lubricant is added completely or in part

-26-

to the mixture. When added in part to the mixture, the remaining part of the lubricant is sprayed on the punches and/or the dies.

According to still another embodiment of the invention all of the lubricant is sprayed
5 on the punches and/or the dies, and the step of adding lubricant to the mixture is then obviously omitted.

Compression of the mixture may be carried out with any compressing apparatus used in
10 the art, e.g. by a rotational compression machine.

The hardness of the tablet is adapted so that the resulting brittleness, as measured by European Pharmacopoeia methods, is less than 2% by weight, preferably 1% by weight.

15 The shape of the tablets may be round, oval, oblong, with a surface that is flat, concave or convex, and optionally may be embossed. In general, the tablets have a mass that ranges from 0.1 to 2.0 grams, and a diameter size ranging from 6 to 18 mm. In case of oblong tablets, the length may vary between 10 and 30 mm, e.g. about 21 mm and the
20 width between 5 and 15 mm e.g. about 10 mm. Preferably the length of the oblong tablets is about double the size of the width. The size and weight of the tablets will be determined by the amount of active ingredient that is incorporated therein.

Examples

Example 1

25 1. Preparation of the particles

Production was carried out on a GPCG1 fluidized bed in Bottom Spray mode.

The spraying solution was prepared by dissolving TMC125 in a mixture of solvents
96% alcohol/methylene with HPMC 2910 5cPs (supplied by Dow Chemical) and
30 polysorbate 20 (Montanox®20, supplied by SEPPIC).

Lactose EFK (supplied by HMS) was introduced on the fluidized bed and the solution
was sprayed in the bottom spray mode. The granules were dried on a fluidized bed and
milled using a FITZMILL grinder with a 790-µm grid .

35 After the spraying step, the resulting granules were dried on the fluidized bed.

Preparation of the spraying solution

Composition of the spraying solutions
40

Raw materials	Quantity (kg)
TMC125	11.67
HPMC 2910 5cPs	35.00
Montanox 20DF	1.17
Methylene chloride	860.00
Ethyl alcohol 96%	83.33
Total	991.17

Six batches of the spraying solutions (hereafter referred to as solutions 1 to 6) were made according to the following operating method:

1. Pour the alcohol and methylene chloride in the tank after weighing them in the same container
2. Put under stirring, add TMC125 and wait until complete dissolution
3. Dissolve Montanox 20 DF under stirring
4. Dissolve HPMC 2910 cPs under stirring until it is completely dissolved
5. Readjust the solution weight by adding methylene chloride to make up for the evaporation which occurred during the solution preparation.

Solutions 1 & 2 were prepared at first. Then, solutions 3 & 4 and solutions 5 & 6 were prepared respectively during the spraying of solutions 1 & 2 and solutions 3 & 4.

Layering with solution 1 and 2

1. 63 kg of Lactose EFK were introduced into the fluidized bed GPCG 120 and solutions 1 and 2 were sprayed. The granulates were then dried in the fluidized bed GPCG 120 for approximately fifteen minutes.
2. The granulates were sieved on a 1.60 mm screen after this first layering step. The undersized granulates were milled half part on a Fitzmill grinder equipped with a 790 μ m screen and half part on a Forplex grinder to optimize the process time.
3. The oversized granulates were milled on a Fitzmill grinder equipped with a 790 μ m screen.
4. The resulting mass of granulates was then divided into two equivalent parts: Part A and Part B. Each part was composed of half of granulates milled with a Forplex grinder and half of granulates milled with a Fitzmill grinder to have an homogeneous particle size distribution.

Layering with solutions 3 and 4

These two solutions were sprayed on part A. The resulting granulates were dried in the fluidized bed GPCG 120 for approximately one hour. They were then sieved on a 1.60 mm screen and the oversized were milled on a Fitzmill grinder equipped with a 790 μ m screen and / or on a Forplex grinder.

Layering with solutions 5 and 6

These two solutions were sprayed on part B. The resulting granulates were dried in the fluidized bed GPCG 1120 for approximately one hour and sieved on a 1.60 mm screen. The oversized were milled on a Fitzmill grinder equipped with a 790 μ m screen and / or on a Forplex grinder.

Spraying rate for solutions 1 and 2 was set at about 3.00 kg/min, for solutions 3 and for (sprayed on part A) about 3.25 kg/min and for solutions 5 and 6 (sprayed on part B) also about 3.25 kg/min. Spraying pressure was 3.5 bar.

Drying Step

Parts A and B were dried at 80°C for about 72 hours.

Milling Step

The dry granulates were milled on a Forplex grinder. One cycle was sufficient to reach the desired particle size (between 150 to 300 μ m).

Final blend

Blend all the milled granulates for 12 minutes at 10 rpm.

Total Composition

Raw materials	Quantity (kg)	Percentage (%)
TMC125	70	20
Lactose EFK	63	18
HPMC 2910 5 cPs	210	60
Montanox 20 DF	7	2
Methylene chloride	5160	/
Ethyl alcohol 96%	500	/
Total	350	100

The resulting quantity of granulates was about 350 kg, yield about 95%.

2. Results

2.1 Granulometric distribution

The granulometric distribution of various batches obtained by the above described procedures were in the following ranges :

5

D_{10%}: 58-94 µm

D_{50%}: 229-267 µm

D_{90%}: 471-496 µm

10 Example 2: Preparation of Tablets

Excipients	Quantity (mg)	Percentage (%)
TMC125 granulates	1000.00	76.92
Microcrystalline cellulose PH302*	212.90	16.38
Reticulated Polyvidone CL USP	72.80	5.60
Silica USP (Aerosil)**	3.90	0.30
Hydrogenated cotton oil**	10.40	0.80
Total	1300.00	100.00

* The quantity of Microcrystalline Cellulose is adjusted according to the actual assay of TMC125 granulates to have a tablet with a constant weight of 1300 mg.

15 ** An external lubrication of micronised hydrogenated cotton oil + 0.9% Aerosil is performed to solve sticking phenomenon. The quantity sprayed and fixed is not included in the tablet formula (not quantifiable).

Manufacturing of tablets

20 Blend

The blend was performed in a 1500 l container with a Soneco blender:

1. Weight and sieve Reticulated Polyvidone CL + Microcrystalline Cellulose PH 302 + Anhydrous colloidal Silica 200 on a 630 µm screen.
Blend then for 12 minutes at 10 rpm.
- 25 2. Add TMS 125 granulates and blend for 12 minutes at 10 rpm.
3. Add the hydrogenated cotton oil after sieving on a 630µm screen and blend for 1 minute at 10 rpm.
4. The blend was sampled according to the analytical protocol but no analytical tests were performed on the blend.

30

-30-

Tableting

For the manufacture of these four tablets batches, the tablet press used was the industrial Fette P1200.

Oblong punches (21.59 x 10.29 mm) were used.

- 5 An external lubrication of micronised hydrogenated cotton oil + 0.9% of Aerosil was applied on the punches and the dies to avoid sticking.

CLAIMS

1. A process for preparing a particle comprising a co-precipitate surrounding a neutral hydrophilic carrier, said process comprising spraying an organic solution on a neutral hydrophilic carrier, said solution comprising at least one triazine or pyrimidine active ingredient having HIV inhibiting properties, one surface active agent, and one hydrophilic polymer, wherein the spraying of whole of the solution occurs in at least two separate steps, each of these steps followed by a grinding step of the product obtained at the end of the preceding step.
2. The process for the preparation of particles according to Claim 1, comprising the following steps:
 - a) preparing a solution comprising at least one triazine or pyrimidine active ingredient having HIV inhibiting properties, a hydrophilic polymer, and a surface active agent, in an organic solvent;
 - b) spraying of part of the solution obtained in step a) on a neutral hydrophilic carrier;
 - c) grinding of the particles obtained in step b);
 - d) spraying of the remaining quantity of organic solution on the particles obtained in step c), and
 - e) final grinding of the particles obtained in step d).
3. The process for the preparation of particles according to any one of claims 1 and 2, wherein the spraying/grinding sequence (steps b through d) is repeated once, or several times.
4. The process for the preparation of particles according to one of any of the claims 1 to 3, wherein procedures additionally comprise a drying step, either after each spraying step before grinding, or immediately following grinding.
5. The process for the preparation of particles according to any one of the claims 1 to 4, wherein the inert hydrophilic carrier consists of any chemically or pharmaceutically inert excipient, existing in a particle form, crystalline or amorphous, and selected preferably from the group consisting of sugars, sugar derivatives, celluloses, or mixtures thereof.
6. The process for the preparation of particles according to any one of the claims 1 to 5, wherein the hydrophilic polymer is selected from the group of

polyvinylpyrrolidones, and in particular polymers with a molecular weight ranging from 10,000 to 50,000, cellulose derivatives, preferably hydroxypropylmethylcellulose, hydroxypropylcellulose, hydroxymethylcellulose, hydroxypropylmethylcellulose phthalate and hydroxypropylmethylcellulose aceto-
5 succinate, acrylic polymers and polyethylene glycols.

7. The process for the preparation of particles according to any one of the claims 1 to 6, wherein the surface active agent is selected in the group consisting of cationic, anionic, non-ionic or amphoteric agents, separately or as a mixture thereof.
10
8. The process for the preparation of particles according to any one of the claims 1 to 7, wherein the organic solvent is selected in the group consisting of ethanol, isopropanol, tetrahydrofuran, isopropyl ether, acetone, methylethylacetone, methylene chloride, and mixtures of these solvents.
15
9. The process for the preparation of particles according to any one of the claims 1 to 8, wherein the spraying steps are carried out in a coating pan, in a perforated pan, or on a fluidized bed.
- 20 10. A particle comprising a co-precipitate applied in a layer surrounding a neutral hydrophilic carrier, and comprising at least one antiviral pyrimidine and triazine, at least one surface-active agent, and at least one hydrophilic polymer.
- 25 11. The particle according to claim 10 obtainable by the process as claimed in any of claims 1 – 9.
12. The particle according to claims 10 - 11, wherein the surface-active agent is present in the particle in an amount that varies from 1 to 60 % by weight.
- 30 13. The particle according to any one of the Claims 10 and 12, wherein the inert hydrophilic carrier is present in an amount of up to 95% by weight.
14. The particle according to any one of the Claims 10 to 13, wherein the weight ratio of the hydrophilic polymer to the active ingredient ranges from 10:1 to 1:2.
35
15. The particle according to any one of the Claims 10 to 14, herein the surface-active agent is present in an amount that may vary from 0.1 to 20 % in weight compared to the total resulting mass.

16. The particle according to any one of the Claims 10 to 15, wherein the unit particle
size of the inert hydrophilic carrier ranges from 50 μm to 500 μm , preferably from 90
5 μm to 200 μm .

17. A pharmaceutical dosage form, wherein said form contains at least one particle
according to any one of the claims 10 to 15, optionally combined with
pharmaceutically acceptable excipients.
10

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/53 A61P31/18 A61K9/16 A61K9/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, EMBASE, MEDLINE, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
Y	WO 00/27825 A (AKEN KOEN JEANNE ALFONS VAN ; KOYMANS LUCIEN MARIA HENRICUS (BE); HEER) 18 May 2000 (2000-05-18) cited in the application page 20, line 30 - page 21, line 3	1-17
Y	EP 1 103 252 A (LICONSA LIBERACION CONTROLADA) 30 May 2001 (2001-05-30) claim 1	1-17
A	US 5 776 495 A (DUCLOS ROSELYNE ET AL) 7 July 1998 (1998-07-07) claims 1,2,8,9	1-17

☐ Further documents are listed in the continuation of box C☒ Patent family members are listed in annex

* Special categories of cited documents

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

G document member of the same patent family

Date of the actual completion of the international search

13 January 2005

Date of mailing of the international search report

24/01/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Sindel, U

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0027825	A	18-05-2000	AT 233740 T 15-03-2003
			AU 762523 B2 26-06-2003
			AU 6200899 A 29-05-2000
			BG 105418 A 30-11-2001
			BR 9915552 A 14-08-2001
			CA 2350801 A1 18-05-2000
			CN 1322198 T 14-11-2001
			CZ 20011533 A3 17-10-2001
			DE 69905683 D1 10-04-2003
			DE 69905683 T2 18-03-2004
			DK 1002795 T3 30-06-2003
			EA 4049 B1 25-12-2003
			EE 200100252 A 15-10-2002
			WO 0027825 A1 18-05-2000
			EP 1270560 A1 02-01-2003
			EP 1002795 A1 24-05-2000
			ES 2193664 T3 01-11-2003
			HK 1025330 A1 13-06-2003
			HR 20010161 A1 28-02-2002
			HU 0104177 A2 28-03-2002
			ID 28376 A 17-05-2001
			JP 2002529456 T 10-09-2002
			NO 20011696 A 04-04-2001
			NZ 511116 A 29-08-2003
			PL 347586 A1 08-04-2002
			PT 1002795 T 31-07-2003
			SI 1002795 T1 31-10-2003
			SK 6032001 A3 07-01-2002
			TR 200101306 T2 22-10-2001
			US 2003114472 A1 19-06-2003
			US 2004039005 A1 26-02-2004
			ZA 200103769 A 12-08-2002
EP 1103252	A	30-05-2001	ES 2157731 A1 16-08-2001
			AT 234077 T 15-03-2003
			AU 750730 B2 25-07-2002
			AU 4782199 A 14-02-2000
			BG 105282 A 30-11-2001
			BR 9912300 A 17-04-2001
			CA 2337509 A1 03-02-2000
			DE 69905905 D1 17-04-2003
			DE 69905905 T2 04-03-2004
			DK 1103252 T3 14-07-2003
			EE 200100041 A 17-06-2002
			EP 1103252 A1 30-05-2001
			HK 1037531 A1 18-07-2003
			HR 20010128 A1 31-12-2002
			HU 0103708 A2 28-02-2002
			IL 140910 A 28-03-2004
			JP 2002521323 T 16-07-2002
			NO 20010289 A 07-03-2001
			NZ 509886 A 27-09-2002
			PL 345625 A1 02-01-2002
			SI 1103252 T1 31-10-2003
			SK 862001 A3 06-08-2001
			US 6737082 B1 18-05-2004
			CZ 20010175 A3 16-05-2001
			WO 0004881 A1 03-02-2000

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/051545

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
EP 1103252	A	ES 2194480 T3	16-11-2003	
		PT 1103252 T	31-07-2003	
		RU 2209628 C2	10-08-2003	
		TR 200100177 T2	23-07-2001	
		ZA 200101293 A	21-08-2001	
<hr/>				
US 5776495	A	07-07-1998	WO 9704749 A1	13-02-1997
			FR 2722984 A1	02-02-1996
			AU 3082995 A	26-02-1997
			EP 0761208 A1	12-03-1997
			FI 962978 A	20-01-1997
			JP 10505574 T	02-06-1998
<hr/>				